

2021-11

# Tephra deposition and bonding with reactive oxides enhances burial of organic carbon in the Bering Sea

Longman, J

<http://hdl.handle.net/10026.1/18129>

---

10.1029/2021GB007140

Global Biogeochemical Cycles: an international journal of global change

American Geophysical Union

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# Tephra deposition and bonding with reactive oxides enhances burial of organic carbon in the Bering Sea

Jack Longman<sup>1, 2, \*</sup> (ORCID: 0000-0002-2725-2617), Thomas M. Gernon<sup>2</sup> (ORCID: 0000-0002-7717-2092), Martin R. Palmer<sup>2</sup>, Hayley R. Manners<sup>2,3</sup> (ORCID: 0000-0001-8545-4463).

<sup>1</sup>Marine Isotope Geochemistry, Institute for Chemistry and Biology of the Marine Environment (ICBM), University of Oldenburg, 26129 Oldenburg, Germany

<sup>2</sup>School of Ocean & Earth Science, University of Southampton, Southampton SO14 3ZH, UK

<sup>3</sup>School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, PL4 8AA, United Kingdom

\* Corresponding author: Jack Longman: jack.longman@uni-oldenburg.de

## Key Points

- Tephra layers are loci of marine organic carbon (OC) burial with distinct carbon isotopic compositions.
- Preservation primarily linked to association of OC with reactive iron phases, accounting for ~80% of all OC in tephra layers.
- Distribution of reactive iron from tephra into surrounding sediment has enhanced OC burial in these layers (~33% of OC in sediments bound to reactive phases).
- OC-reactive Fe coupling is observed in sediments >700,000 years old, indicating long-term persistence of these complexes.
- These interactions may help explain how OC is preserved in sediments on geological timescales.

## Abstract

Preservation of organic carbon (OC) in marine sediments exerts a major control on the cycling of carbon in the Earth system. In these marine environments, OC preservation may be enhanced by diagenetic reactions in locations where deposition of fragmental volcanic material called tephra occurs. While the mechanisms by which this process occurs are well understood, site-specific studies of this process are limited. Here, we report a study of sediments from the Bering Sea (IODP Site U1339D) to investigate the effects of marine tephra deposition on carbon cycling during the Pleistocene and Holocene. Our results suggest that tephra layers are loci of OC burial with distinct  $\delta^{13}\text{C}$  values, and that this process is primarily linked to bonding of OC with reactive metals, accounting for ~80% of all OC within tephra layers. In addition, distribution of reactive metals from the tephra into non-volcanic sediments above and below the tephra layers enhances OC preservation in these sediments, with ~33% of OC bound to reactive phases. Importantly, OC-Fe coupling is evident in sediments >700,000 years old. Thus, these interactions may help explain the observed preservation of OC in ancient marine sediments.

## Plain Language Summary

The burial of organic carbon in marine sediments is one of the major carbon sinks on Earth, meaning that it removes carbon dioxide from the ocean-atmosphere system. However, the speed at which burial occurs varies across the globe, and is dependent on a range of factors, from the amount of nutrients in the water column, to the type of sediment. Despite evidence suggesting that when tephra is deposited to the seafloor carbon burial is enhanced, very little work has been done to investigate this process. We have therefore analyzed sediments from the Bering Sea, where volcanoes from the Aleutian Islands and Kamchatka regularly deposit tephra in the ocean. We found that organic carbon burial is indeed associated with ash deposition, and importantly, that organic carbon is preserved in the ash layers themselves. We show here that this carbon is preserved effectively because of chemical reactions between the organic carbon and reactive iron, which is released by the ash, creating conditions which preserve carbon for hundreds of thousands of years.

## Introduction

The preservation of organic carbon (OC) in marine sediments exerts a controlling influence on the carbon cycle, providing a link between the active pools (e.g. oceans, atmosphere and terrestrial environments) and the inactive, long-term carbon pools, such as those within sedimentary rocks (Arndt et al., 2013; Burdige, 2007; LaRowe et al., 2020). Only about 0.5% of all organic matter produced in the oceans is ultimately preserved in the sedimentary record, with the remainder remineralised and reintroduced into active carbon pools (Hedges & Keil, 1995). The process of OC burial and preservation leads to net removal of  $\text{CO}_2$  from the atmosphere, thus any process which changes the marine sedimentary carbon sink is critically important for understanding the climate system on geological timescales.

Explosive volcanism delivers approximately  $1 \text{ km}^3$  of ash to the atmosphere every year (Pyle, 1995), and because a high proportion of volcanoes are located close to the oceans, and 70% of the world is covered by oceans, much of this material falls into seawater and onto seafloor substrates (Olgun et al., 2011; Pyle, 1995). Tephra also enters the oceans through rapid erosion of newly created volcanic deposits (Cashman et al., 2013). This material eventually settles to the seafloor, and is deposited in the sedimentary record as tephra layers (Dingwell et al., 2012; Pyle, 1989). Tephra may also derive

from submarine eruptions, such that tephra is thought to represent as much as 25% of marine sediments in the Pacific Ocean (Scudder et al., 2009; Straub & Schmincke, 1998).

There are four mechanisms by which enhanced preservation of OC in marine sediments may occur as a result of tephra deposition and diagenesis: (1) fertilization; (2) reactive metal bonding; (3) reduced oxidant exposure, and (4) authigenic carbonate formation (Longman et al., 2019, 2020). Upon deposition in the ocean, and as a result of the dissolution of reactive mineral phases, tephra releases large amounts of macro- and micronutrients such as P, Fe and Mn (Frogner et al., 2001; Jones & Gislason, 2008) that may alleviate deficiencies (Moore et al., 2013), particularly when Fe is the limiting nutrient. Indeed, the fertilization effect has been observed in the form of phytoplankton blooms in the aftermath of tephra deposition events (Achterberg et al., 2013; Duggen et al., 2010; Langmann et al., 2010; Uematsu et al., 2004). Tephra deposition on the seafloor rapidly reduces pore water O<sub>2</sub> contents to near-zero as a result of oxidation of silicate-bound Fe<sup>II</sup> (Haeckel et al., 2001; Hembury et al., 2012), thus inhibiting the oxidation and remineralization of OC.

Reactive metal bonding is thought to account for ~20% of all OC preserved in marine sediments (Lalonde et al., 2012), and reactive Fe, Mn and Al phases are known to be released from tephra layers to adjacent sediments during diagenesis (Homoky et al., 2011). Hence, while it has yet to be directly observed, tephra deposition likely also contributes enhanced OC preservation through this process.

The association of abundant tephra layers and high OC concentrations has been taken to suggest that tephra diagenesis played a role in enhanced OC preservation in ancient environments (Lee et al., 2018; Tang et al., 2020), and there is growing evidence of tephra-related processes actively enhancing OC preservation in modern sediments (Hembury et al., 2012; Homoky et al., 2011; Murray et al., 2018). Nevertheless, the role of reactive metals released by tephra in this process has not been studied in detail.

Here, we estimate the contribution of this process, and the others outlined above, to the preservation of OC in the Bering Sea. Our work investigates the changing chemistry above, below and within, tephra layers deposited throughout the Quaternary Period (2.6 Ma to present), helping to improve our understanding of the impact tephra deposition has on marine sedimentary organic carbon.

## Methods and Materials

### Study Site

Sediments from IODP Expedition 323 site U1339D (54°31.26'N, 169°44.35'W, 200 mbsl), on the Umnak Plateau in the Bering Sea (Fig. 1) largely comprise two lithological end-members: biogenic diatom-rich sediment and a tephra. The volcanogenic material is sourced from eruptions along the Aleutian arc, and constitutes ~4 – 40% of the sediment (Takahashi et al., 2011; Vaughn & Caissie, 2017). Tephra layers were identified on-ship by their difference in colour (typically black) with respect to the adjacent sediments, abrupt lower and upper contacts, and the presence of glass shards in smear slides (Takahashi et al., 2011). After visual identification of cores, subsampling of both tephra layers and background sediment was undertaken in this study. The surface of the split-core sections was removed and care was taken to ensure that only the centre of the tephra layers was sampled to reduce contamination from the adjacent sediments below visibly detectable levels. In all studied examples, a sharp boundary between background sediment and the base of the tephra layer is present (e.g., Figs. S1-S2). Mixing of sediment into the top of the tephra is present in some tephra layers,

but this is clearly apparent in the core and, where present, was typically limited to the uppermost 2-3 cm of tephra layers (Figs. S1-S2).

For this study, we selected 9 sections, which are denoted Sections 1 to 9, with their depths and approximate ages indicated in Figure 2 (see Table S1 for details). Indicative ages were taken from the biostratigraphic age model of Takahashi et al. (2011b), interpolating linearly between the midpoint of each datum (see Fig. 2, Table S1).

## Geochemical Analyses

### Organic Carbon

Organic carbon measurements were carried out on a Vario PYRO cube Element Analyser (EA) coupled to a vision isotope ratio mass spectrometer (IRMS) at the University of Southampton. Approximately 20 mg of homogenised sample was acidified in perchloric acid to remove any carbonate prior to multiple rinses with Milli-Q water. EA quality control was performed via repeated measurements of High Organic Sediment Standard (HOSS; Element Microanalysis Ltd., n=11), with a reproducibility of  $\pm 0.07$  wt %, and Acetalinide (n=8), with a reproducibility of  $\pm 0.1$  wt % (1 SD).

Bulk sediment carbon isotope signatures ( $\delta^{13}\text{C}_{\text{Bulk}}$ ) were measured on  $\text{CO}_2$  evolved from EA combustion, and calibrated to USGS 40 and USGS 41a, with reproducibility of  $\pm 0.02$  ‰ (n=5), and  $\pm 0.36$  ‰ (n=4), respectively (1 SD). Repeat analyzes of HOSS (n=11), and Acetanilide (n=8) were used for quality control, with precision of  $\pm 0.04$  ‰ and  $\pm 0.05$  ‰ respectively (1 SD).

### Inorganic Carbon

Approximately 20 mg of homogenised sediment was analyzed via coulometry of perchloric acid treatment-released  $\text{CO}_2$ , using an AutoMate Prep Device (AutoMate FX, Inc., Bushnell, Florida, USA) using a UIC CM5015  $\text{CO}_2$  Coulometer (UIC Inc., Joliet, Illinois, USA) at the University of Southampton. Calibration was performed using a pure carbonate standard (CAS #471-34-1), and quality control was completed via analysis of an in-house stream sediment standard.

Subsamples of layers which contained quantifiable levels of  $\text{CaCO}_3$  were then selected for carbonate carbon and oxygen isotope analysis. According to the  $\text{CaCO}_3$  content, between 5-15 mg of sample was analysed via a Thermo Scientific Kiel IV Carbonate device coupled to a MAT253 IRMS at the University of Southampton. Perchloric acid released  $\text{CO}_2$ , which was analysed for carbon and oxygen isotopes ( $\delta^{13}\text{C}_{\text{Carb}}$  and  $\delta^{18}\text{O}$ ). Replicate analyses of an in-house standard were calibrated to NBS-18 and NBS-19, with reproducibility of  $\pm 0.13$  ‰ and  $\pm 0.12$  ‰ for  $\delta^{13}\text{C}_{\text{Carb}}$ , and  $\pm 0.17$  ‰ and  $\pm 0.23$  ‰ for  $\delta^{18}\text{O}$  (all 1 SD).

### Elemental Geochemistry

Bulk sample geochemistry was carried out after digestion at  $130^\circ\text{C}$  for 24 hours via a closed-vessel mixed acid ( $\text{HNO}_3$ - $\text{HCl}$ - $\text{HF}$ ) approach. Digests were then diluted to 2%  $\text{HNO}_3$  and analyzed on a Thermo Scientific X-Series ICP-MS at the University of Southampton. Here, we present data for Al, Mn, Fe and Ba (Table S1). Alongside samples, blanks and reference material (HISS-1 and JMS-1 marine sediment standards) were prepared and analyzed in the same manner (see Table S2 for HISS-1 recoveries and blank values).

### Reactive Oxides

To isolate reactive oxide phases, a 4 hour dithionite extraction was performed (Kostka & Luther, 1994; Lalonde et al., 2012; Mehra & Jackson, 1958). Despite the potential drawbacks of this approach, which include incomplete extraction of all OC bound to reactive phases (Fisher et al., 2020), it remains the most widely utilised method for investigating interactions between reactive phases and OC (Faust et al., 2021; Fisher et al., 2021).

For each sample, 4 ml of dithionite reagent (buffered to pH 4.8) was prepared, and added to 0.1 g of homogenised, freeze-dried sediment. To maintain pH 4.8, a buffered 0.35M sodium acetate, 0.2M sodium citrate solution was used, and heated to 60°C in a water bath. Samples were agitated using a vortex mixer every 15 minutes. This approach has been previously used to extract amorphous Fe-oxides alongside a fraction of crystalline Fe-oxides and acid volatile sulphides (Kostka & Luther, 1994; Roy et al., 2013). Dithionite-extracted fractions were diluted in the same manner as for elemental analysis and analyzed on a Thermo Scientific X-Series ICP-MS at the University of Southampton. Results are presented in Table S3. Chilean Margin sediment (RR9702A-42MC, see Muratli et al., (2012)), was prepared and analysed in the same manner as the samples, with results for reactive Fe ( $Fe_R$ ) and reactive Mn ( $Mn_R$ ) found to be within the range of previously reported values (Table S2). For  $Fe_R$ , values of  $10475 \pm 125$  ppm (1 SD,  $n=3$ ) are close to previously measured values of  $10800 \pm 800$  ppm (Roy et al., 2013) and  $9300 \pm 200$  ppm (N.A. Murray et al., 2016). For  $Mn_R$ , measured values of  $306 \pm 15$  ppm (1SD,  $n=3$ ) compare well with other studies, including  $290 \pm 10$  ppm (Murray et al., 2016), and  $300 \pm 60$  ppm (Roy et al., 2013) (Table S2).

To investigate the composition of carbon associated with the phases extracted via dithionite leaching, we used the approach of Lalonde et al. (2012). This involves analysis of the OC content before and after the extraction experiment outlined above, and analysis of  $\delta^{13}C_{bulk}$  before and after extraction. In addition, a control experiment was completed, where samples were extracted using sodium chloride instead of sodium dithionite and trisodium citrate, according to the method of Lalonde et al., (2012). For tephra, this released 0.004 wt% of the OC, and for sediment 0.02 wt%. These values were then used to correct experimental data (Lalonde et al., 2012; Shields et al., 2016), although it has been shown this approach can result in underestimations of  $Fe_R$ -associated OC (Fisher et al., 2020) (Table S4). For simplicity, and in a similar manner to previous studies (e.g. Faust et al., 2021), we consider the results of the extraction experiment to represent  $Fe_R$ -bound OC, and not  $Mn_R$ - and  $Al_R$ -bound OC. Results are presented in Table S4.

Using the results of the extraction experiment and control experiment, the fraction of OC associated with reactive phases (hereafter  $f_{OC-Fe}$ ) and the isotopic composition of this OC (hereafter  $\delta^{13}C_{Fe-OC}$ ) were calculated using the following equation (Lalonde et al., 2012):

$$f_{Fe-OC} = \frac{OC_{extract} - OC_{control}}{OC_{bulk}}$$

where  $f_{OC-Fe}$  is the fraction of OC bound to reactive phases,  $OC_{control}$  is the OC content after control extraction (for either tephra or sediment),  $OC_{extract}$  is OC content after dithionite extraction and  $OC_{bulk}$  is OC content prior to extraction. Using the precision data derived from the EA standards ( $\pm 0.07$  wt %), and the average wt % OC content of the samples lost during the extraction (0.32 wt%), we estimate the error on  $f_{OC-Fe}$  to be on average  $\pm 21\%$ . Only two samples display an absolute OC loss lower than the analytical error (Table S3). The isotopic composition of the fraction of OC extracted ( $\delta^{13}C_{Fe-OC}$ ) was calculated using the following equation:

$$\delta^{13}C_{Fe-OC} = \frac{\delta^{13}C_{bulk} \times OC_{bulk} - \delta^{13}C_{Fe-OC-extract} \times OC_{extract}}{OC_{bulk} - OC_{extract}}$$

where  $\delta^{13}\text{C}_{\text{bulk}}$  is the  $\delta^{13}\text{C}$  of OC before the dithionite experiment and  $\delta^{13}\text{C}_{\text{Fe-OC-extract}}$  is the  $\delta^{13}\text{C}$  of OC after the dithionite extraction. Using the calculated  $\delta^{13}\text{C}_{\text{Fe-OC}}$ , and the absolute amount of OC associated with  $\text{Fe}_R$  ( $\text{OC}_{\text{Fe}}$ ), the isotopic composition of the non  $\text{Fe}_R$ -bound OC ( $\delta^{13}\text{C}_{\text{Non-Fe-OC}}$ ) was then calculated using the following equation:

$$\delta^{13}\text{C}_{\text{Non-Fe-OC}} = \frac{\delta^{13}\text{C}_{\text{Fe-OC}} \times \text{OC}_{\text{Fe}} - \delta^{13}\text{C}_{\text{bulk}} \times \text{OC}_{\text{bulk}}}{\text{OC}_{\text{Fe}} - \text{OC}_{\text{bulk}}}$$

Using a similar approach to that outlined above, precision from IRMS standards ( $\pm 0.13\text{‰}$ ), the average  $\delta^{13}\text{C}_{\text{Fe-OC}}$  ( $0.56\text{‰}$ ), we calculate error on  $\delta^{13}\text{C}_{\text{Fe-OC}}$  estimates to be on average  $\pm 22\%$ . In this case, only one sample displays a  $\delta^{13}\text{C}$  shift lower than analytical error (Table S3).

## Palaeoproductivity

To assess changing palaeoproductivity, we used the biogenic fraction of barium ( $\text{Ba}_{\text{Bio}}$ ), a commonly used proxy (Schoepfer et al., 2015). This approach first calculates the proportion of excess Ba in the sediments, an approach which uses the expected ratio of Ba to the conservative element Al in detrital, non-biogenic Ba to calculate the remainder:

$$\text{Ba}_{\text{Bio}} = \text{Ba}_{\text{Total}} - \text{Al}_{\text{Total}} \times (\text{Ba}/\text{Al}_{\text{detrital}})$$

For  $\text{Ba}/\text{Al}_{\text{detrital}}$ , we assume the primary detrital contributor is tephra and use an average value from all tephra layers in this study (0.0099). Using published biostratigraphic ages (Takahashi, Ravelo, & Alvarez Zarikian, 2011), we calculate accumulation rates which are then used to convert raw  $\text{Ba}_{\text{Bio}}$  into  $\text{Ba}_{\text{Bio}}$  flux:

$$\text{Ba}_{\text{Bio}}\text{Flux} = \text{Ba}_{\text{Bio}} \times \rho \times \text{LSR}$$

where LSR is the linear sedimentation rate, in  $\text{cm}^{-1} \text{kyr}$  and  $\rho$  is the density of sediment, estimated using the following equation:

$$\rho = 0.0794 \times \ln(x) + 0.650$$

where  $x$  is the age of the sample in kyr (Schoepfer et al., 2015). Results may be found in Table S1.

## Results

The composition of the tephra layers and adjacent non-volcanogenic sediments are compared in Table 1. The tephra layers show lower average OC and inorganic carbon contents, but extend to much higher inorganic carbon concentrations in Section 6 (Fig. 2). The bulk  $\delta^{13}\text{C}$  values of the two groups overlap, but the tephra layers have more negative mean  $\delta^{13}\text{C}$  values ( $-25.4 \pm 1.18\text{‰}$ , 1SD,  $n=22$ ) than those of the sediments ( $-23.91 \pm 0.6\text{‰}$ , 1SD,  $n=44$ ) (Table S5). The tephra layers are slightly enriched in total Fe and Mn with similar levels of Al to sediment. The tephra layers contain slightly lower reactive phase contents (Fig. 3; Table 1). The dithionite extraction experiment shows a greater average  $f_{\text{OC-Fe}}$  in tephra layers ( $79 \pm 13\%$ , 1SD,  $n=13$ ) than sediments ( $33 \pm 22\%$ , 1SD,  $n=24$ ), with  $\delta^{13}\text{C}_{\text{Fe-OC}}$  in tephra layers and sediments averaging  $-25.83\text{‰}$  and  $-24.16\text{‰}$ , respectively (Table 1; Fig. 4, Table S4). Carbonate  $\delta^{13}\text{C}$  analyses show two clusters for  $\delta^{13}\text{C}_{\text{Carb}}$  (Fig. 5). One cluster ( $n=5$ ) displays a narrow range in  $\delta^{13}\text{C}_{\text{Carb}}$  values between  $-1.01$  and  $-1.58\text{‰}$ . The other ( $n=8$ ) has  $\delta^{13}\text{C}_{\text{Carb}}$  values between  $-13.41$  and  $-19.56\text{‰}$ .  $\delta^{18}\text{O}$  values also differ between the two clusters, with samples in the first lying between  $-7.09$  and  $1.34\text{‰}$  and the second between  $2.92$  and  $9.2\text{‰}$  (Fig. 5).

## Discussion

## 221 OC in tephra layers

222 Our analyses show that all tephra layers contain a component of OC, with an average of  $0.3 \pm 0.7$  wt%  
223 (1SD,  $n=22$ ), compared to an average of 0.9 wt% in the background sediments (Fig. 2a, Table S5).  
224 Because fresh tephra contains negligible OC, these data indicate that some OC preservation  
225 mechanism occurred within the tephra layers. Bulk carbon isotope analyses of the tephra layers and  
226 surrounding sediments indicate that the composition of this OC is different to what is preserved in  
227 surrounding sediments. Mean tephra  $\delta^{13}\text{C}$  is  $-25.4 \pm 1.2\text{‰}$  (1SD,  $n=22$ ) and mean sediment  $\delta^{13}\text{C}$  is  $-$   
228  $23.9 \pm 0.6\text{‰}$  (1SD,  $n=44$ ) (Fig. 2d), with statistically different means and variances (T-test p-value  
229  $<0.005$ ). This suggests that the tephra layers contain a distinct source of  $^{13}\text{C}$ -depleted carbon, which  
230 is consistent with the fact that we minimized sampling of any background sedimentary OC mixed  
231 into the tephra layers, despite the bioturbated nature of some sediments at site U1339D. However,  
232 where present, the effect of bioturbation on tephra is limited to the uppermost 1-2 cm of the layers  
233 (Takahashi et al., 2011). Therefore, our sampling method—which avoided gradational boundaries  
234 and sampled the centre of the tephras—deliberately avoided the sampling of any potentially mixed  
235 sediment (Figs. S1-2). Further, if bioturbation were to have played a role, one would expect the  
236 tephra layers to contain a mix of OC brought in from surrounding layers, rather than a distinct source  
237 as we observe. It is possible the distinct  $\delta^{13}\text{C}$  represents a shift to OC of a more terrestrial origin in  
238 tephras, with Yukon River OC typically on the order of  $-27\text{‰}$  (Guo & Macdonald, 2006). However,  
239 this is unlikely to constitute a significant proportion of the OC supply to our site, with the Yukon  
240 delta located nearly 1000 km to the northeast. In addition, there is no a priori reason to expect the  
241 tephra layers to contain more terrestrial OC than the sediments, but the possibility cannot be  
242 discounted.

243 An additional explanation for the distinct OC  $\delta^{13}\text{C}$  values in the tephra may be autochthonous  
244 microbial biomass formation, because microbial fatty acid  $\delta^{13}\text{C}$  values range from  $-30$  to  $-45\text{‰}$ ,  
245 depending on the carbon source (Cifuentes & Salata, 2001; Gong & Hollander, 1997; Hayes, 2001).  
246 This occurs due to fractionation processes which take place as microbes utilise carbon, the majority  
247 of which result in  $\delta^{13}\text{C}$  depletion (Hayes, 2001). While we cannot be certain of a biomass origin, the  
248 negative values of microbial biomass means that only a relatively small contribution is necessary to  
249 result in the observed isotopic shift. Further study would be required to confirm this hypothesis, but  
250 evidence suggests that volcanic glass may provide the ideal substrate for microbial growth (Li et al.,  
251 2020; Zhang et al., 2017). Indeed, tephra layers have been shown to contain microbial communities  
252 which are distinct, more diverse and greater in number than surrounding sediments (Inagaki et al.,  
253 2003), with sulfide oxidation suggested as an energy source (Böhnke et al., 2019). However, even if  
254 microbial biomass plays a role in the OC preserved in these tephras, both the lack of direct evidence,  
255 and the relatively un-depleted  $\delta^{13}\text{C}$  values in tephra suggest that another source of OC is also  
256 present within the tephra layers. One further possibility is that the shift in lithology from sediment to  
257 tephra has allowed for the preferential preservation of certain organic compounds, due to changing  
258 reactivity and chemical compositions. This could be linked to the preservation of OC via bonding  
259 with reactive iron ( $\text{Fe}_\text{R}$ ) phases, as a result of the high  $\text{Fe}_\text{R}$  content in ash (Homoky et al., 2011), and  
260 there is evidence to suggest that Fe is released during tephra alteration (Luo et al., 2020; Maters et  
261 al., 2017). Such a mechanism is not dependent on the formation mechanism of the OC, rather  
262 increasing preservation across all OC forms, as the complexes formed are difficult to break down  
263 (Lalonde et al., 2012). This may result in the depletion of  $\delta^{13}\text{C}$  in the tephras, as OC degradation  
264 tends to result in the retained OC containing a more negative  $\delta^{13}\text{C}$  value (Lehmann et al., 2002;  
265 Zonneveld et al., 2010), but further research is necessary to confirm this. As such, it may be that the  
266 OC retained in the tephra is simply marine OC which has undergone selective preservation, resulting



in more depleted  $\delta^{13}\text{C}$  than surrounding sediment (Fig. 4). Thus, a microbial component cannot be discounted, but also cannot be proven from our data alone.

## Reactive metal bonding

Globally, ~20% of marine OC is thought to be preserved via bonding with  $\text{Fe}_\text{R}$  phases (Barber et al., 2017; Lalonde et al., 2012), hence we have investigated how bonding reactions may have influenced OC preservation at U1339D. Absolute values of OC associated with Fe (OC-Fe) are similar for both tephra (average 0.31 wt %) and sediment (average 0.27 wt %), but with much more variability in the tephra layers (Fig. 4c; Figure S3). However, the variability in the bulk OC content, and the lower OC content of tephra, means that the percentage of OC associated with  $\text{Fe}_\text{R}$  ( $f_{\text{OC-Fe}}$ ) is clearly different between the tephra and background sediment (Fig. 4e, Table 1). The  $f_{\text{OC-Fe}}$  in the background sediment (average 33%) is similar to the global average, but within the tephra layers,  $f_{\text{OC-Fe}}$  increases to an average of 79% (Fig. 4; Table 1). By comparison, the highest  $f_{\text{OC-Fe}}$  observed in marine sediments elsewhere is ~40% in deltaic sediments and ~30% in sediments underlying the equatorial Pacific upwelling zone (Lalonde et al., 2012). The intense  $\text{Fe}_\text{R}$ -OC bonding within the tephra layers may be linked to the high proportion of  $\text{Fe}^\text{II}$  within tephra deposits (Homoky et al., 2011), that provides an ideal environment for OC inner-sphere bonding (Barber et al., 2017). Interestingly, the  $\text{Fe}_\text{R}$  content of the tephra layers at Site U1339C is lower than those in the background sediments (Fig. 3), potentially because Aleutian eruptions are primarily andesitic and rhyolitic as opposed to basaltic in composition (Figure S4). However,  $f_{\text{OC-Fe}}$  is high, supporting recent work which suggests the absolute availability of  $\text{Fe}_\text{R}$  is not the dominant control on OC-Fe association (Faust et al., 2021). The higher proportion of  $f_{\text{OC-Fe}}$  in the sediments adjacent to the tephra (relative to the more distal sediments) (Fig. 4) may be related to the diffusion of colloidal reactive Fe out of the tephra layers (cf. Homoky et al., 2011).

The isotopic signature of the  $\text{Fe}_\text{R}$ -bound OC ( $\delta^{13}\text{C}_{\text{Fe-OC}}$ ) may indicate the type of OC being preserved via these interactions (Fig 4b, d, f). As with bulk  $\delta^{13}\text{C}$ , the  $\delta^{13}\text{C}_{\text{Fe-OC}}$  of tephra layers is consistently more negative than those in the sediments, with an average of -25.50 ‰ in tephra, and -22.4 ‰ in sediments (Fig. 4f), suggesting a distinct carbon source. The affinity of marine OC to  $\text{Fe}_\text{R}$  phases has been observed in a range of marine sediments located on the continental shelf previously, likely indicating a marine source of sediment-hosted OC bound to  $\text{Fe}_\text{R}$  (Lalonde et al., 2012), a finding in line with those from other marine environments such as estuaries (Sirois et al., 2018; Zhao et al., 2018). As with bulk sediment OC, it is possible that the more negative nature of  $\delta^{13}\text{C}_{\text{Fe-OC}}$  in tephra represents a shift toward microbial OC generation and preservation (Cifuentes & Salata, 2001; Gong & Hollander, 1997), resulting from fractionation which occurs as microbes utilise carbon (Hayes, 2001). Circumstantial evidence for this hypothesis comes from laboratory studies which demonstrate that reactive Fe oxides may act as electron suppliers for metabolism of metal-reducing bacteria (Coker et al., 2012; Kato et al., 2010).

However, as with bulk  $\delta^{13}\text{C}$  values, the variation may result from the preservation of certain OC compounds. Our data appear to support this, as the bulk  $\delta^{13}\text{C}$  value of tephra is similar to that of tephra-hosted  $\delta^{13}\text{C}_{\text{Fe-OC}}$  values (Fig. 4), a function of the majority of OC being bound to  $\text{Fe}_\text{R}$  phases in tephra. The retention of a depleted signal in  $\delta^{13}\text{C}_{\text{Fe-OC}}$  suggests that what remains in the tephra is the non-labile,  $\text{Fe}_\text{R}$ -complexed OC, and that the loss of labile compounds has caused the depletion. Indeed, in background sediments,  $\delta^{13}\text{C}_{\text{Fe-OC}}$  displays less negative isotope ratios (Fig. 4). This suggests that outside of tephra layers the isotopic composition has not been shifted as Fe-OC makes up a smaller proportion of total OC and so less preservation of negative  $\delta^{13}\text{C}$  is occurring. The

enhancement of OC preservation due to  $\text{Fe}_R$  bonding in tephra layers may thus provide a previously unconsidered sink for such OC in sediments containing abundant tephra (Hedges et al., 1997).

The molar ratio of organic carbon to reactive iron (OC:Fe; Fig. 5a, Table S4) may provide information on the mechanisms of binding between OC and Fe (Faust et al., 2021; Lalonde et al., 2012), with low ratios indicative of simple mono-layer sorption, and higher ratios related to coprecipitation (Wagai & Mayer, 2007). In the Bering Sea sections, OC:Fe molar ratios vary greatly, suggesting a range of OC-Fe interactions, but tephra layers typically display lower OC:Fe than in the surrounding sediments (Fig. 5a). The high OC:Fe ratios observed in some layers (OC:Fe >10) may indicate deposition under anoxic/sub-oxic conditions (Lalonde et al., 2012). In tephra layers, this is likely related to the consumption of porewater  $\text{O}_2$  during tephra diagenesis (Hembury et al., 2012). Low OC:Fe ratios are typically linked to  $\text{O}_2$  exposure (Lalonde et al., 2012), or terrestrial OC-Fe bonding (Barber et al., 2014; Faust et al., 2021), but as discussed above these mechanisms are unlikely to be at play here. If microbial activity is a contributor to tephra OC, utilisation of the reactive Fe during microbial metabolism may have altered the  $\text{Fe}_R$  content adsorbed or coprecipitated to other OC (Elizabeth Cooper et al., 2017; Eusterhues et al., 2014). Further, since the dithionite extraction removes all “reactive” Fe phases, and not simply those complexed with OC, it is possible that low OC:Fe ratios are related to the extraction of  $\text{Fe}_R$  phases not involved with bonding (Faust et al., 2021). By pairing our OC:Fe data with  $f_{\text{OC-Fe}}$ , and comparing with previous studies, it is clear that the OC- $\text{Fe}_R$  interaction in tephra layers differ from any previous studies (Fig. 6). Our sediment data are close to previous studies of oxic and suboxic sediments (Lalonde et al., 2012), but the tephra show very high  $f_{\text{OC-Fe}}$  associated with low OC:Fe. This further indicates that a process not typically associated with marine sediments occurs in the tephra layers (Fig. 6).

### Long term persistence of an enhanced ‘rusty carbon sink’ in tephra-rich sediments

Regardless of the source of the OC preserved in the sediments and tephra, there is evidence of extensive OC- $\text{Fe}_R$  bonding in all layers of the studied Bering Sea sediments, with particularly high levels of OC complexation in both the tephra and surrounding sediments (Fig. 4). These values suggest that the environment produced by tephra deposition, in which enhanced availability of nutrients (and potentially  $\text{Fe}_R$  although not in the Bering Sea) is coupled to localised oxygen depletion (Hembury et al., 2012), is one in which high proportions of local OC are complexed to  $\text{Fe}_R$ . The enhanced  $f_{\text{OC-Fe}}$  proportions in tephra are greater than reported in any surface sediments, even in anoxic depositional environments, suggesting the size of the ‘rusty sink’ in any given sediment may be more related to the availability of  $\text{Fe}_R$  phases and not the available oxygen (Lalonde et al., 2012).

In addition, even in the oldest layers around the tephra deposited between 700-745 kyr (i.e., section 21H4; Takahashi et al., 2011), all  $f_{\text{OC-Fe}}$  proportions are above 10%, and greater than 70% in the tephra. This suggests that bonding with reactive oxides provides a long-term sink for OC in the marine environment, one which persists for far longer than previously indicated (Faust et al., 2021). This has implications for long-term carbon cycling on Earth, suggesting OC-Fe bonding may represent an important component of the high activation energy ( $E$ ) OC involved in the long-term preservation of marine OC (Hemingway et al., 2019). The proportion of high- $E$  organic compounds has been shown to increase as time proceeds in marine sediments, with our work suggesting a proportion of this is linked to OC- $\text{Fe}_R$  bonds. This finding, coupled with the implication that tephra are loci of intense OC- $\text{Fe}_R$  bonding, suggest that tephra may be involved in the burial and long-term

sequestration of OC after periods of major volcanic activity, such as in the Late Ordovician (Buggisch et al., 2010) and the mid-Cretaceous (Lee et al., 2018).

## Other processes involved in OC preservation

Another potential carbon sink in tephra is authigenic carbonate (Schrag et al., 2013), which may form in tephra layers themselves (Longman et al., 2021), or sediments in which levels of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  have been enhanced by ash deposition (Hong et al., 2020; Longman et al., 2019; Luo et al., 2020; Torres et al., 2020). In most tephra and sediments at site U1139D, there is little evidence for this process occurring (Fig. 2b), potentially due to the small amount of carbonate precipitation typically promoted by ash alteration (Hong et al., 2020), but there are exceptions, particularly in one layer where carbonate contents exceed 50 wt% (tephra 7; Fig. 2b). In addition, a small number of sediment layers show carbonate enrichment (Fig. 2b).

Carbon and oxygen isotope analyses of the tephra-hosted carbonates ( $\delta^{13}\text{C}_{\text{carb}}$ ) indicate that there are two clear groups (Fig. 5b). The first, composed solely of carbonate from sediment layers, is characterized by  $\delta^{13}\text{C}_{\text{carb}}$  between -1 and -2 ‰, and appears to be indicative of a biogenic carbonate formation, or authigenic carbonates formed from dissolved inorganic carbon in seawater (e.g. Humphreys et al., 2015). The second group shows  $\delta^{13}\text{C}_{\text{carb}}$  values between -12 to -20 ‰ (Fig. 5b). This  $^{12}\text{C}$  depletion is typical of carbonates formed as a result of the anaerobic oxidation of methane once this methane reaches the zone where it occurs in concert with sulfate (Sivan et al., 2007; Whiticar & Faber, 1986). Authigenic carbonates ( $\text{C}_{\text{auth}}$ ) formed as a result of this process may act as a carbon sink, preventing the methane from returning carbon to the ocean, and locking it into stable carbonate phases (Schrag et al., 2013). Previous work in the region has shown widespread evidence for  $\text{C}_{\text{auth}}$  in Bering sediment (Pierre et al., 2016), with similarly  $^{12}\text{C}$ -depleted carbon isotope signatures (Hein et al., 1979). As we see little evidence for  $\text{C}_{\text{auth}}$  formation in sediments, and with previous work showing  $\text{C}_{\text{auth}}$  in tephra layers (Hein et al., 1979), it is possible that tephra alteration has supplied the  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  necessary for formation. If true, this would suggest  $\text{C}_{\text{auth}}$  formation is a carbon sink enhanced by tephra diagenesis.

It is also possible that tephra deposition may stimulate phytoplankton productivity in surface seawater (Langmann et al., 2010; Olgun et al., 2011). Using both OC content and biogenic barium flux ( $\text{Ba}_{\text{Bio}}$ ) as proxies for palaeoproductivity (Schoepfer et al., 2015), we investigated the impact of tephra deposition on productivity in the Bering Sea. There is little evidence of increased productivity in sediments directly surrounding tephra deposits, with slightly lower average OC content, and similar  $\text{Ba}_{\text{Bio}}$  (Fig. 2c). This is despite evidence of plankton blooms in the aftermath of eruptions in the region (Hamme et al., 2010; Langmann et al., 2010). This suggests either that: i) plankton blooms are transitory and short-lived, having very little impact on overall productivity in the region; or ii) the organic carbon produced by such blooms is either not exported from the upper ocean prior to remineralisation, or it is transported to other locations by ocean currents. In addition, the Bering Sea is typically an area of high productivity (Wehrmann et al., 2011), so that the addition of tephra makes little difference to overall production. A final possibility is that the andesitic and relatively low-Fe nature of the tephra deposited in the Bering Sea means that it does not contain sufficient amounts of nutrient to influence biological productivity.

## Conclusions

Our results demonstrate enhanced OC preservation in tephra, and in the sediments surrounding tephra deposits. The OC in the tephra layers is primarily associated with reactive metal phases, with an average  $f_{\text{OC-Fe}}$  value of 77% in tephra. Thus, tephra layers contain the highest  $f_{\text{OC-Fe}}$  proportions yet

reported. Isotopic analyses indicate that this OC is primarily marine in origin, supporting previous studies which demonstrate the affiliation of Fe<sub>R</sub> to marine OC in marine environments (Sirois et al., 2018; Zhao et al., 2018). The data also shows the viability of long-term stability of such relationships, with high f<sub>OC-Fe</sub> proportions in sediments older than 700 kyr. This finding may explain observed increases in OC activation energy as age increases (Hemingway et al., 2019), with OC-Fe<sub>R</sub> interactions resulting in hard to break down organic compounds. In addition to bonding of OC with reactive metal phases, there is limited evidence for authigenic carbonate formation in these tephra layers.

## References

- Achterberg, E. P., Moore, C. M., Henson, S. A., Steigenberger, S., Stohl, A., Eckhardt, S., et al. (2013). Natural iron fertilization by the Eyjafjallajökull volcanic eruption. *Geophysical Research Letters*, 40(5), 921–926. <https://doi.org/10.1002/grl.50221>
- Arndt, S., Jørgensen, B. B., LaRowe, D. E., Middelburg, J. J., Pancost, R. D., & Regnier, P. (2013). Quantifying the degradation of organic matter in marine sediments: A review and synthesis. *Earth-Science Reviews*, 123, 53–86. <https://doi.org/10.1016/j.EARSCIREV.2013.02.008>
- Barber, A., Lalonde, K., Mucci, A., & Gélinas, Y. (2014). The role of iron in the diagenesis of organic carbon and nitrogen in sediments: A long-term incubation experiment. *Marine Chemistry*, 162, 1–9. <https://doi.org/10.1016/j.marchem.2014.02.007>
- Barber, A., Brandes, J., Leri, A., Lalonde, K., Balind, K., Wirick, S., et al. (2017). Preservation of organic matter in marine sediments by inner-sphere interactions with reactive iron. *Scientific Reports*, 7(1), 366. <https://doi.org/10.1038/s41598-017-00494-0>
- Böhnke, S., Sass, K., Gonnella, G., Diehl, A., Kleint, C., Bach, W., et al. (2019). Parameters Governing the Community Structure and Element Turnover in Kermadec Volcanic Ash and Hydrothermal Fluids as Monitored by Inorganic Electron Donor Consumption, Autotrophic CO<sub>2</sub> Fixation and 16S Tags of the Transcriptome in Incubation Experiments. *Frontiers in Microbiology*, 10, 2296. <https://doi.org/10.3389/FMICB.2019.02296>
- Buggisch, W., Joachimski, M. M., Lehnert, O., Bergström, S. M., Repetski, J. E., & Webers, G. F. (2010). Did intense volcanism trigger the first Late Ordovician icehouse? *Geology*, 38(4), 327–330. <https://doi.org/10.1130/G30577.1>
- Burdige, D. J. (2007). Preservation of organic matter in marine sediments: Controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chemical Reviews*, 107(2), 467–485. <https://doi.org/10.1021/cr050347q>
- Cashman, K. V., Stephen, R., & Sparks, J. (2013). How volcanoes work: A 25 year perspective. *Bulletin of the Geological Society of America*, 125(5–6), 664–690. <https://doi.org/10.1130/B30720.1>
- Cifuentes, L. A., & Salata, G. G. (2001). Significance of carbon isotope discrimination between bulk carbon and extracted phospholipid fatty acids in selected terrestrial and marine environments. *Organic Geochemistry*, 32(4), 613–621. [https://doi.org/10.1016/S0146-6380\(00\)00198-4](https://doi.org/10.1016/S0146-6380(00)00198-4)
- Coker, V. S., Byrne, J. M., Telling, N. D., Van Der Laan, G., Lloyd, J. R., Hitchcock, A. P., et al. (2012). Characterisation of the dissimilatory reduction of Fe(III)-oxyhydroxide at the microbe - mineral interface: The application of STXM-XMCD. *Geobiology*, 10(4), 347–354. <https://doi.org/10.1111/j.1472-4669.2012.00329.x>
- Dingwell, D. B., Lavallée, Y., & Kueppers, U. (2012). Volcanic ash: A primary agent in the Earth

440 system. *Physics and Chemistry of the Earth, Parts A/B/C*, 45–46, 2–4.  
 441 <https://doi.org/10.1016/J.PCE.2011.07.007>

442 Duggen, S., Olgun, N., Croot, P., Hoffmann, L., Dietze, H., Delmelle, P., & Teschner, C. (2010). The role  
 443 of airborne volcanic ash for the surface ocean biogeochemical iron-cycle: a review.  
 444 *Biogeosciences*, 7(3), 827–844. <https://doi.org/10.5194/bg-7-827-2010>

445 Elizabeth Cooper, R., Eusterhues, K., Wegner, C. E., Uwe Totsche, K., & Küsel, K. (2017). Ferrihydrite-  
 446 associated organic matter (OM) stimulates reduction by *Shewanella oneidensis* MR-1 and a  
 447 complex microbial consortia. *Biogeosciences*, 14(22), 5171–5188. [https://doi.org/10.5194/bg-](https://doi.org/10.5194/bg-14-5171-2017)  
 448 14-5171-2017

449 Eusterhues, K., Hädrich, A., Neidhardt, J., Küsel, K., Keller, T. F., Jandt, K. D., & Totsche, K. U. (2014).  
 450 Reduction of ferrihydrite with adsorbed and coprecipitated organic matter: Microbial reduction  
 451 by *Geobacter bremensis* vs. abiotic reduction by Na-dithionite. *Biogeosciences*, 11(18), 4953–  
 452 4966. <https://doi.org/10.5194/bg-11-4953-2014>

453 Faust, J. C., Tessin, A., Fisher, B. J., Zindorf, M., Papadaki, S., Hendry, K. R., et al. (2021). Millennial  
 454 scale persistence of organic carbon bound to iron in Arctic marine sediments. *Nature*  
 455 *Communications*, 12(1), 1–9. <https://doi.org/10.1038/s41467-020-20550-0>

456 Fisher, B. J., Moore, O. W., Faust, J. C., Peacock, C. L., & März, C. (2020). Experimental evaluation of  
 457 the extractability of iron bound organic carbon in sediments as a function of carboxyl content.  
 458 *Chemical Geology*, 556, 119853. <https://doi.org/10.1016/j.chemgeo.2020.119853>

459 Fisher, B. J., Faust, J. C., Moore, O. W., Peacock, C. L., & März, C. (2021). Technical note: Uncovering  
 460 the influence of methodological variations on the extractability of iron-bound organic carbon.  
 461 *Biogeosciences*, 18(11), 3409–3419. <https://doi.org/10.5194/BG-18-3409-2021>

462 Fisk, M. R., Giovannoni, S. J., & Thorseth, I. H. (1998). Alteration of oceanic volcanic glass: Textural  
 463 evidence of microbial activity. *Science*, 281(5379), 978–980.  
 464 <https://doi.org/10.1126/science.281.5379.978>

465 Frogner, P., Reynir Gíslason, S., & Óskarsson, N. (2001). Fertilizing potential of volcanic ash in ocean  
 466 surface water. *Geology*, 29(6), 487. [https://doi.org/10.1130/0091-](https://doi.org/10.1130/0091-7613(2001)029<0487:FPOVAI>2.0.CO;2)  
 467 7613(2001)029<0487:FPOVAI>2.0.CO;2

468 Gong, C., & Hollander, D. J. (1997). Differential contribution of bacteria to sedimentary organic  
 469 matter in oxic and anoxic environments, Santa Monica Basin, California. In *Organic*  
 470 *Geochemistry* (Vol. 26, pp. 545–563). Pergamon. [https://doi.org/10.1016/S0146-](https://doi.org/10.1016/S0146-6380(97)00018-1)  
 471 6380(97)00018-1

472 Hamme, R. C., Webley, P. W., Crawford, W. R., Whitney, F. A., DeGrandpre, M. D., Emerson, S. R., et  
 473 al. (2010). Volcanic ash fuels anomalous plankton bloom in subarctic northeast Pacific.  
 474 *Geophysical Research Letters*, 37(19), n/a-n/a. <https://doi.org/10.1029/2010GL044629>

475 Hayes, J. M. (2001). Fractionation of carbon and hydrogen isotopes in biosynthetic processes.  
 476 *Reviews in Mineralogy and Geochemistry*, 43(1), 191–277.  
 477 <https://doi.org/10.2138/gsrmg.43.1.225>

478 Hedges, J. I., Keil, R. G., & Benner, R. (1997). What happens to terrestrial organic matter in the  
 479 ocean? In *Organic Geochemistry* (Vol. 27, pp. 195–212). Pergamon.  
 480 [https://doi.org/10.1016/S0146-6380\(97\)00066-1](https://doi.org/10.1016/S0146-6380(97)00066-1)

481 Hedges, John I., & Keil, R. G. (1995). Sedimentary organic matter preservation: an assessment and  
 482 speculative synthesis. *Marine Chemistry*, 49(2–3), 81–115. [https://doi.org/10.1016/0304-](https://doi.org/10.1016/0304-4203(95)00008-F)  
 483 4203(95)00008-F

484 Hein, J. R., O’Neil, J. R., Jones, O’NEIL, J. R., & JONES, M. G. (1979). Origin of authigenic carbonates in  
 485 sediment from the deep Bering Sea. *Sedimentology*, 26(5), 681–705.  
 486 <https://doi.org/10.1111/j.1365-3091.1979.tb00937.x>

487 Hembury, D. J., Palmer, M. R., Fones, G. R., Mills, R. A., Marsh, R., & Jones, M. T. (2012). Uptake of  
 488 dissolved oxygen during marine diagenesis of fresh volcanic material. *Geochimica et*  
 489 *Cosmochimica Acta*, 84, 353–368. <https://doi.org/10.1016/J.GCA.2012.01.017>

490 Hemingway, J. D., Rothman, D. H., Grant, K. E., Rosengard, S. Z., Eglinton, T. I., Derry, L. A., & Galy, V.  
 491 V. (2019). Mineral protection regulates long-term global preservation of natural organic  
 492 carbon. *Nature*, 570(7760), 228–231. <https://doi.org/10.1038/s41586-019-1280-6>

493 Homoky, W. B., Hembury, D. J., Hepburn, L. E., Mills, R. A., Statham, P. J., Fones, G. R., & Palmer, M.  
 494 R. (2011). Iron and manganese diagenesis in deep sea volcanogenic sediments and the origins  
 495 of pore water colloids. *Geochimica et Cosmochimica Acta*, 75(17), 5032–5048.  
 496 <https://doi.org/10.1016/J.GCA.2011.06.019>

497 Hong, W. L., Torres, M. E., & Kutterolf, S. (2020). Towards a global quantification of volcanogenic  
 498 aluminosilicate alteration rates through the mass balance of strontium in marine sediments.  
 499 *Chemical Geology*, 550, 119743. <https://doi.org/10.1016/j.chemgeo.2020.119743>

500 Humphreys, M. P., Achterberg, E. P., Griffiths, A. M., McDonald, A., & Boyce, A. J. (2015).  
 501 Measurements of the stable carbon isotope composition of dissolved inorganic carbon in the  
 502 northeastern Atlantic and Nordic Seas during summer 2012. *Earth System Science Data*, 7(1),  
 503 127–135. <https://doi.org/10.5194/essd-7-127-2015>

504 Inagaki, F., Suzuki, M., Takai, K., Oida, H., Sakamoto, T., Aoki, K., et al. (2003). Microbial Communities  
 505 Associated with Geological Horizons in Coastal Subseafloor Sediments from the Sea of Okhotsk.  
 506 *Applied and Environmental Microbiology*, 69(12), 7224–7235.  
 507 <https://doi.org/10.1128/AEM.69.12.7224-7235.2003>

508 Jones, M. T., & Gislason, S. R. (2008). Rapid releases of metal salts and nutrients following the  
 509 deposition of volcanic ash into aqueous environments. *Geochimica et Cosmochimica Acta*,  
 510 72(15), 3661–3680. <https://doi.org/10.1016/j.gca.2008.05.030>

511 Kato, S., Nakamura, R., Kai, F., Watanabe, K., & Hashimoto, K. (2010). Respiratory interactions of soil  
 512 bacteria with (semi)conductive iron-oxide minerals. *Environmental Microbiology*, 12(12), 3114–  
 513 3123. <https://doi.org/10.1111/j.1462-2920.2010.02284.x>

514 Kostka, J. E., & Luther, G. W. (1994). Partitioning and speciation of solid phase iron in saltmarsh  
 515 sediments. *Geochimica et Cosmochimica Acta*, 58(7), 1701–1710.  
 516 [https://doi.org/10.1016/0016-7037\(94\)90531-2](https://doi.org/10.1016/0016-7037(94)90531-2)

517 Lalonde, K., Mucci, A., Ouellet, A., & Gélinas, Y. (2012). Preservation of organic matter in sediments  
 518 promoted by iron. *Nature*, 483(7388), 198–200. <https://doi.org/10.1038/nature10855>

519 Langmann, B., Zakšek, K., Hort, M., & Duggen, S. (2010). Volcanic ash as fertiliser for the surface  
 520 ocean. *Atmos. Chem. Phys. Atmospheric Chemistry and Physics*, 10, 3891–3899. Retrieved from  
 521 [www.atmos-chem-phys.net/10/3891/2010/](http://www.atmos-chem-phys.net/10/3891/2010/)

522 LaRowe, D. E., Arndt, S., Bradley, J. A., Estes, E. R., Hoarfrost, A., Lang, S. Q., et al. (2020, May 1). The  
523 fate of organic carbon in marine sediments - New insights from recent data and analysis. *Earth-*  
524 *Science Reviews*. Elsevier B.V. <https://doi.org/10.1016/j.earscirev.2020.103146>

525 Lee, C.-T. A., Jiang, H., Ronay, E., Minisini, D., Stiles, J., & Neal, M. (2018). Volcanic ash as a driver of  
526 enhanced organic carbon burial in the Cretaceous. *Scientific Reports*, 8(1), 4197.  
527 <https://doi.org/10.1038/s41598-018-22576-3>

528 Lehmann, M. F., Bernasconi, S. M., Barbieri, A., & McKenzie, J. A. (2002). Preservation of organic  
529 matter and alteration of its carbon and nitrogen isotope composition during simulated and in  
530 situ early sedimentary diagenesis. *Geochimica et Cosmochimica Acta*, 66(20), 3573–3584.  
531 [https://doi.org/10.1016/S0016-7037\(02\)00968-7](https://doi.org/10.1016/S0016-7037(02)00968-7)

532 Li, L., Bai, S., Li, J., Wang, S., Tang, L., Dasgupta, S., et al. (2020). Volcanic ash inputs enhance the  
533 deep-sea seabed metal-biogeochemical cycle: A case study in the Yap Trench, western Pacific  
534 Ocean. *Marine Geology*, 430, 106340. <https://doi.org/10.1016/j.margeo.2020.106340>

535 Longman, J., Palmer, M. R., Gernon, T. M., & Manners, H. R. (2019). The role of tephra in enhancing  
536 organic carbon preservation in marine sediments. *Earth-Science Reviews*, 192, 480–490.  
537 <https://doi.org/10.1016/j.earscirev.2019.03.018>

538 Longman, J., Palmer, M. R., & Gernon, T. M. (2020). Viability of greenhouse gas removal via artificial  
539 addition of volcanic ash to the ocean. *Anthropocene*, 32.  
540 <https://doi.org/10.1016/j.ancene.2020.100264>

541 Longman, J., Gernon, T. M., Palmer, M. R., Jones, M. T., Stokke, E. W., & Svensen, H. H. (2021).  
542 Marine diagenesis of tephra aided the Paleocene-Eocene Thermal Maximum termination. *Earth*  
543 *and Planetary Science Letters*, 571, 117101. <https://doi.org/10.1016/J.EPSL.2021.117101>

544 Luo, M., Torres, M. E., Hong, W. L., Pape, T., Fronzek, J., Kutterolf, S., et al. (2020). Impact of iron  
545 release by volcanic ash alteration on carbon cycling in sediments of the northern Hikurangi  
546 margin. *Earth and Planetary Science Letters*, 541, 116288.  
547 <https://doi.org/10.1016/j.epsl.2020.116288>

548 Maters, E. C., Delmelle, P., & Gunnlaugsson, H. P. (2017). Controls on iron mobilisation from volcanic  
549 ash at low pH: Insights from dissolution experiments and Mössbauer spectroscopy. *Chemical*  
550 *Geology*, 449, 73–81. <https://doi.org/10.1016/J.CHEMGEO.2016.11.036>

551 Mehra, O. P., & Jackson, M. L. (1958). Iron Oxide Removal from Soils and Clays by a Dithionite-Citrate  
552 System Buffered with Sodium Bicarbonate. *Clays and Clay Minerals*, 7(1), 317–327.  
553 <https://doi.org/10.1346/CCMN.1958.0070122>

554 Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., et al. (2013).  
555 Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*, 6(9), 701–710.  
556 <https://doi.org/10.1038/ngeo1765>

557 Muratli, J. M., McManus, J., Mix, A., & Chase, Z. (2012). Dissolution of fluoride complexes following  
558 microwave-assisted hydrofluoric acid digestion of marine sediments. *Talanta*, 89, 195–200.  
559 <https://doi.org/10.1016/j.talanta.2011.11.081>

560 Murray, N.A., Muratli, J. M., Hartwell, A. M., Manners, H., Megowan, M. R., Goñi, M., et al. (2016).  
561 Data report: dissolved minor element compositions, sediment major and minor element  
562 concentrations, and reactive iron and manganese data from the Lesser Antilles volcanic arc

563 region, IODP Expedition 340 Sites U1394, U1395, U1396, U1399, and U1400. *Proceedings of the*  
564 *Integrated Ocean Drilling Program, 340*. <https://doi.org/10.2204/iodp.proc.340.207.2016>

565 Murray, Natalie A., McManus, J., Palmer, M. R., Haley, B., & Manners, H. (2018). Diagenesis in  
566 tephra-rich sediments from the Lesser Antilles Volcanic Arc: Pore fluid constraints. *Geochimica*  
567 *et Cosmochimica Acta*, 228, 119–135. <https://doi.org/10.1016/J.GCA.2018.02.039>

568 Norrman, B., Zwiefel, U. L., Hopkinson, C. S., & Brian, F. (1995). Production and utilization of  
569 dissolved organic carbon during an experimental diatom bloom. *Limnology and Oceanography*,  
570 40(5), 898–907. <https://doi.org/10.4319/LO.1995.40.5.0898>

571 Ogawa, H., Amagai, Y., Koike, I., Kaiser, K., & Benner, R. (2001). Production of refractory dissolved  
572 organic matter by bacteria. *Science*, 292(5518), 917–920.  
573 <https://doi.org/10.1126/science.1057627>

574 Olgun, N., Duggen, S., Croot, P. L., Delmelle, P., Dietze, H., Schacht, U., et al. (2011). Surface ocean  
575 iron fertilization: The role of airborne volcanic ash from subduction zone and hot spot  
576 volcanoes and related iron fluxes into the Pacific Ocean. *Global Biogeochemical Cycles*, 25(4),  
577 n/a-n/a. <https://doi.org/10.1029/2009GB003761>

578 Pierre, C., Blanc-Valleron, M. M., Caquineau, S., März, C., Ravelo, A. C., Takahashi, K., & Alvarez  
579 Zarikian, C. (2016). Mineralogical, geochemical and isotopic characterization of authigenic  
580 carbonates from the methane-bearing sediments of the Bering Sea continental margin (IODP  
581 Expedition 323, Sites U1343-U1345). *Deep-Sea Research Part II: Topical Studies in*  
582 *Oceanography*, 125–126, 133–144. <https://doi.org/10.1016/j.dsr2.2014.03.011>

583 Pyle, D. M. (1989). The thickness, volume and grain size of tephra fall deposits. *Bulletin of*  
584 *Volcanology*, 51(1), 1–15. <https://doi.org/10.1007/BF01086757>

585 Pyle, D. M. (1995). Mass and energy budgets of explosive volcanic eruptions. *Geophysical Research*  
586 *Letters*, 22(5), 563–566. <https://doi.org/10.1029/95GL00052>

587 Roy, M., McManus, J., Goñi, M. A., Chase, Z., Borgeld, J. C., Wheatcroft, R. A., et al. (2013). Reactive  
588 iron and manganese distributions in seabed sediments near small mountainous rivers off  
589 Oregon and California (USA). *Continental Shelf Research*, 54, 67–79.  
590 <https://doi.org/10.1016/J.CSR.2012.12.012>

591 Schoepfer, S. D., Shen, J., Wei, H., Tyson, R. V., Ingall, E., & Algeo, T. J. (2015). Total organic carbon,  
592 organic phosphorus, and biogenic barium fluxes as proxies for paleomarine productivity. *Earth-*  
593 *Science Reviews*, 149, 23–52. <https://doi.org/10.1016/J.EARSCIREV.2014.08.017>

594 Schrag, D. P., Higgins, J. A., Macdonald, F. A., & Johnston, D. T. (2013). Authigenic carbonate and the  
595 history of the global carbon cycle. *Science*, 339(6119), 540–3.  
596 <https://doi.org/10.1126/science.1229578>

597 Scudder, R. P., Murray, R. W., & Plank, T. (2009). Dispersed ash in deeply buried sediment from the  
598 northwest Pacific Ocean: An example from the Izu–Bonin arc (ODP Site 1149). *Earth and*  
599 *Planetary Science Letters*, 284(3–4), 639–648. <https://doi.org/10.1016/J.EPSL.2009.05.037>

600 Shields, M. R., Bianchi, T. S., Gélinais, Y., Allison, M. A., & Twilley, R. R. (2016). Enhanced terrestrial  
601 carbon preservation promoted by reactive iron in deltaic sediments. *Geophysical Research*  
602 *Letters*, 43(3), 1149–1157. <https://doi.org/10.1002/2015GL067388>

603 Sirois, M., Couturier, M., Barber, A., Gélinais, Y., & Chaillou, G. (2018). Interactions between iron and



604 organic carbon in a sandy beach subterranean estuary. *Marine Chemistry*.  
605 <https://doi.org/10.1016/J.MARCHEM.2018.02.004>

606 Sivan, O., Schrag, D. P., & Murray, R. W. (2007). Rates of methanogenesis and methanotrophy in  
607 deep-sea sediments. *Geobiology*, 5, 141–151. [https://doi.org/10.1111/j.1472-](https://doi.org/10.1111/j.1472-4669.2007.00098.x)  
608 [4669.2007.00098.x](https://doi.org/10.1111/j.1472-4669.2007.00098.x)

609 Staudigel, H., Furnes, H., McLoughlin, N., Banerjee, N. R., Connell, L. B., & Templeton, A. (2008,  
610 August 1). 3.5 billion years of glass bioalteration: Volcanic rocks as a basis for microbial life?  
611 *Earth-Science Reviews*. Elsevier. <https://doi.org/10.1016/j.earscirev.2008.04.005>

612 Straub, S. M., & Schmincke, H. U. (1998). Evaluating the tephra input into Pacific Ocean sediments:  
613 distribution in space and time. *Geologische Rundschau*, 87(3), 461–476.  
614 <https://doi.org/10.1007/s005310050222>

615 Takahashi, K., Ravelo, A., & Alvarez-Zarikian, C. (2011). *Proceedings of the Integrated Ocean Drilling*  
616 *Program: Expedition Reports: Bering Sea Paleooceanography*. vol. 323. College Station, Texas:  
617 Integrated Ocean Drilling Program Management International, Inc., for the Integrated Ocean  
618 Drilling Program.

619 Takahashi, K., Ravelo, A. C., & Alvarez Zarikian, C. A. (2011). Site U1339. In *Proceedings of the*  
620 *Integrated Ocean Drilling Program Volume 323* (p. 84). Tokyo: Integrated Ocean Drilling  
621 Program Management International, Inc.

622 Tang, L., Song, Y., Jiang, S., Jiang, Z., Li, Z., Yang, Y., et al. (2020). Organic matter accumulation of the  
623 Wufeng-Longmaxi shales in southern Sichuan Basin: Evidence and insight from volcanism.  
624 *Marine and Petroleum Geology*, 120, 104564.  
625 <https://doi.org/10.1016/j.marpetgeo.2020.104564>

626 Thorseth, I. H., Furnes, H., & Tumyr, O. (1995). Textural and chemical effects of bacterial activity on  
627 basaltic glass: an experimental approach. *Chemical Geology*, 119(1–4), 139–160.  
628 [https://doi.org/10.1016/0009-2541\(94\)00098-5](https://doi.org/10.1016/0009-2541(94)00098-5)

629 Thorseth, I. H., Torsvik, T., Torsvik, V., Daae, F. L., & Pedersen, R. B. (2001). Diversity of life in ocean  
630 floor basalt. *Earth and Planetary Science Letters*, 194(1–2), 31–37.  
631 [https://doi.org/10.1016/S0012-821X\(01\)00537-4](https://doi.org/10.1016/S0012-821X(01)00537-4)

632 Torres, M. E., Hong, W. L., Solomon, E. A., Milliken, K., Kim, J. H., Sample, J. C., et al. (2020, January  
633 1). Silicate weathering in anoxic marine sediment as a requirement for authigenic carbonate  
634 burial. *Earth-Science Reviews*. Elsevier B.V. <https://doi.org/10.1016/j.earscirev.2019.102960>

635 Uematsu, M., Toratani, M., Kajino, M., Narita, Y., Senga, Y., & Kimoto, T. (2004). Enhancement of  
636 primary productivity in the western North Pacific caused by the eruption of the Miyake-jima  
637 Volcano. *Geophysical Research Letters*, 31(6), n/a-n/a. <https://doi.org/10.1029/2003gl018790>

638 Vaughn, D. R., & Caissie, B. E. (2017). Effects of sea-level, sea-ice extent, and nutrient availability on  
639 primary production at the Umnak Plateau, Bering Sea (IODP Site U1339) during Marine Isotope  
640 Stage (MIS) 5. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 485, 283–292.  
641 <https://doi.org/10.1016/J.PALAEO.2017.06.020>

642 Wagai, R., & Mayer, L. M. (2007). Sorptive stabilization of organic matter in soils by hydrous iron  
643 oxides. *Geochimica et Cosmochimica Acta*, 71(1), 25–35.  
644 <https://doi.org/10.1016/j.gca.2006.08.047>

- Wallmann, K., Aloisi, G., Haeckel, M., Tishchenko, P., Pavlova, G., Greinert, J., et al. (2008). Silicate weathering in anoxic marine sediments. *Geochimica et Cosmochimica Acta*, 72(12), 2895–2918. <https://doi.org/10.1016/j.gca.2008.03.026>
- Wehrmann, L. M., Risgaard-Petersen, N., Schrum, H. N., Walsh, E. A., Huh, Y., Ikehara, M., et al. (2011). Coupled organic and inorganic carbon cycling in the deep subseafloor sediment of the northeastern Bering Sea Slope (IODP Exp. 323). *Chemical Geology*, 284(3–4), 251–261. <https://doi.org/10.1016/j.chemgeo.2011.03.002>
- Whiticar, M. J., & Faber, E. (1986). Methane oxidation in sediment and water column environments—Isotope evidence. *Organic Geochemistry*, 10(4–6), 759–768. [https://doi.org/10.1016/S0146-6380\(86\)80013-4](https://doi.org/10.1016/S0146-6380(86)80013-4)
- Zhang, R., Jiang, T., Tian, Y., Xie, S., Zhou, L., Li, Q., & Jiao, N. (2017). Volcanic ash stimulates growth of marine autotrophic and heterotrophic microorganisms. *Geology*, 45(8), G38833.1. <https://doi.org/10.1130/G38833.1>
- Zhao, B., Yao, P., Bianchi, T. S., Shields, M. R., Cui, X. Q., Zhang, X. W., et al. (2018). The Role of Reactive Iron in the Preservation of Terrestrial Organic Carbon in Estuarine Sediments. *Journal of Geophysical Research: Biogeosciences*, 123(12), 3556–3569. <https://doi.org/10.1029/2018JG004649>
- Zonneveld, K. A. F., Versteegh, G. J. M., Kasten, S., Eglinton, T. I., Emeis, K. C., Huguet, C., et al. (2010). Selective preservation of organic matter in marine environments; Processes and impact on the sedimentary record. *Biogeosciences*, 7(2), 483–511. <https://doi.org/10.5194/BG-7-483-2010>

## Acknowledgements

This work was funded by a Natural Environment Research Council (NERC) grant, NE/K00543X/1, “The role of marine diagenesis of tephra in the carbon cycle”. T.M.G. acknowledges support from NERC grant, NE/R004978/1. Datasets for this research are available in the Supplementary Information, and on the Pangaea data archive (doi to follow once assigned)

## Figures




Figure 1: Map of part of the Aleutian Island Arc showing the location of IODP core U1339D. The inset map shows the location of the sampling site within Alaska. Also shown are a number of volcanoes which have actively supplied ash to the Bering Sea during the Quaternary Period (as defined by the Global Volcanism Program of the Smithsonian Institution). Marked in yellow are the key ocean currents affecting sedimentation at this site.





Figure 2: Geochemical parameters of tephra and background sediment from U1339D. a) Organic carbon content in tephra (purple) and background sediments (blue). b)  $\text{CaCO}_3$  content, c) biogenic barium flux, d) bulk  $\delta^{13}\text{C}$ , with average value for both sample type indicated by solid lines. To the left of boxes a-d are the section numbers, depths in metres below sea floor (mbsf) and indicative ages in thousands of years before present (ka). Panels e-h display box and whisker diagrams of the data presented in panels a-d. Boxes are defined between the first and third quartile (interquartile range; IQR), with minimum and maximum whiskers representative of 1.5 times the IQR, and with any outliers ( $>1.5$  times IQR) removed.



690

691 Figure 3: Comparison of total metal contents, reactive metal contents and organic carbon (OC)  
 692 content of sediments and tephras from Site U1339D. Panels a-c display reactive metal  
 693 concentrations plotted against OC whilst panels d-f show total elemental content for Fe, Mn and Al  
 694 against respective reactive metal content. Tephra samples are coloured blue, with sediments in red.




Figure 4: Results of dithionite extraction experiments. In all panels, tephra are indicated by blue circles and background sediments are in red. a) Organic carbon (OC) content in all samples before

extraction versus OC content after extraction. Thick black line indicates where samples should plot if no OC was extracted. Labelled dashed lines indicate the fraction of OC associated with reactive phases ( $f_{OC-Fe}$ ). b) Plot of bulk  $\delta^{13}C$  before and after extraction. Thick black line indicates where samples should plot if no isotopic change were observed. Panels c-d display the same data as a and b but indicate the shift from the original sample (filled circles) to extracted samples (open circles). e)  $f_{OC-Fe}$ , with thick lines indicating average  $f_{OC-Fe}$  for tephra (blue) and sediment (red). f) Bulk isotopic composition of OC associated with reactive phases  $\delta^{13}C_{OC-Fe}$ . As before, thick coloured lines indicate the average  $\delta^{13}C_{OC-Fe}$  of tephra (blue) and sediments (red). For panels c-f, section numbers are indicated to the left. These refer to the ages and depths indicated in Figure 2.




Figure 5: Geochemistry of tephras and sediments from Site U1339D. a) Ratio of OC to reactive Fe, using molar masses for sediments (red) and tephras (blue). b)  $\delta^{13}C$  and  $\delta^{18}O$  of the carbonate fraction (see Methods), with typical values for isotopic composition of seawater (orange rectangle) and biogenic carbonate (yellow rectangle) highlighted. For panel a, section numbers are indicated to the left. These refer to the ages and depths indicated in Figure 2.




Figure 6: Comparison of the fraction of OC associated with FeR ( $f_{OC-Fe}$ ) with the Fe:OC molar ratio from this study and a selection of previous studies. Data from this current study from tephra (blue circles) and sediments (red circles) are shown alongside sediment data from the Barents Sea (green diamonds; Faust et al., 2021). Also shown are data from the compilation of Lalonde et al., (2012), with samples from anoxic/suboxic environments (magenta squares), sulfidic sediments (yellow plus symbols), oxic shelf sediments (orange triangles), deltaic and estuarine sediments (brown stars) and deep-sea sediments (cyan crosses).

Tables

Table 1: Summary of experimental results. Mean, minimum and maximum values for each of the measured variable are presented, for both tephra and sediment layers.

<u>Tephra Layers</u>			
	<u>Mean</u>	<u>Minimum</u>	<u>Maximum</u>
<u>Before dithionite extraction</u>			
Organic carbon content (wt%)	0.33	0.03	3.03



Inorganic carbon content (wt%)	3.02	0	51.66
Ba <sub>Bio</sub> flux (mg cm <sup>-3</sup> kyr)	201	0	2682
Total Fe (wt%)	4.8	2.38	9.58
Total Mn (ppm)	1082	608	2232
Total Al (wt%)	7.98	3.02	13.4
Bulk $\delta^{13}\text{C}$ (‰)	-25.4	-23.23	-27.12
<u>After dithionite extraction</u>			
Organic carbon content (wt%)	0.12	0.01	0.83
Reactive Fe (wt %)	0.57	0.32	1.06
Reactive Mn (ppm)	175	19.83	1042
Reactive Al (ppm)	524.92	287.8	972.8
Bulk $\delta^{13}\text{C}$ (‰)	-26.29	-23.99	-27.96
f <sub>OC-Fe</sub> (%)	0.79	0.55	0.95
$\delta^{13}\text{C}_{\text{Fe-OC}}$ (%)	-25.83	-23.39	-24.16

### Sediment Layers

	<u>Mean</u>	<u>Minimum</u>	<u>Maximum</u>
<u>Before dithionite extraction</u>			
Organic carbon content (wt%)	0.84	0.15	1.41
Inorganic carbon content (wt%)	0.33	0	3.86
Ba <sub>Bio</sub> flux (mg cm <sup>-3</sup> kyr)	166	0	1071
Total Fe (wt%)	4.8	2.88	6.6
Total Mn (ppm)	784.8	488.4	2337
Total Al (wt%)	8.04	6.47	10.03
Bulk $\delta^{13}\text{C}$ (‰)	-23.82	-22.65	-24.71
<u>After dithionite extraction</u>			
Organic carbon content (wt%)	0.61	0.07	1.07
Reactive Fe (wt %)	0.98	0.17	1.87
Reactive Mn (ppm)	170.8	34.32	580.5
Reactive Al (ppm)	914.42	214.4	2254
Bulk $\delta^{13}\text{C}$ (‰)	-24.37	-23.99	-27.96
f <sub>OC-Fe</sub> (%)	0.33	0.06	0.78
$\delta^{13}\text{C}_{\text{Fe-OC}}$ (%)	-24.16	-22.94	-26.39